

SOME RESUPINATE BASIDIOMYCETES FOUND IN PENNSYLVANIA

DANIEL A. DE FIGIO
Illinois State University, Normal

ABSTRACT.—Eleven species of resupinate Basidiomycetes are reported for Pennsylvania for the first time.

Recent collections of fungi previously unreported for Pennsylvania are noted here. The collections were made by the author (D.A.D.) on July 2nd, 1967 from a mixed hardwood forest with a good littor of decaying, deciduous wood. The collection site is located five miles west of Ellitsville, Pennsylvania in Fayette County. All the collections are deposited in the mycological herbarium of Illinois State University (ISU).

Peniophora cremea (Bres.) Sacc. & Syd.

On decayed deciduous wood. Very common, but previously unreported for Pennsylvania. Also known from New York, Ohio, New Jersey, West Virginia, Virginia, Illinois, Alabama, Michigan, Louisiana, Massachusetts, Vermont. VII. 2. 1967. (D.A.D.) 1105, 1106, 1108, 1109, 1111, 1112, 1138, 1143, 1144, 1145, 1070, 1077. (All at ISU).

Peniophora angusta Rog. & Weres.

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from Ontario, British Columbia, Oregon, Rhode Island, and Massachusetts. VII. 2. 1967. (D.A.D.) 1116, 1104, 1069. (All at ISU).

Peniophora aspera (Pers.) Sacc.

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from Illinois, Australia, Denmark, Ukraine, Virginia, and New York. VII. 2. 1967. (D.A.D.) 1115, 1076, 1062. (All at ISU).

Peniophora subulutacea (Karst) Hoehn & Litsch.

On decayed deciduous wood. Previ-

ously unreported for Pennsylvania. Also known from England, France, Finland, Denmark, Poland, New Zealand, Sweden, Austria, British Columbia, Ontario, Illinois, Idaho, Louisiana, Alabama, New York, New Jersey, Maryland, and Washington. VII. 2. 1967. (D.A.D.) 1071. (At ISU).

Peniophora gracillima (Ellis & Everh. ex Rogers & Jackson) G. H. Cunn.

On decayed deciduous wood. Common, but previously unreported for Pennsylvania. Also known from South Africa, Manitoba, New Zealand, Italy, Austria, Australia, England, British Columbia, Ontario, Quebec, Sweden, New York, New Jersey, Vermont, Illinois, New Hampshire, Oregon, and Washington. VII. 2. 1967. (D.A.D.) 1103, 1129, 1130, 1131, 1134. (All at ISU).

Peniophora tenuis (Pat.) Massee

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from Ontario, Austria, Bermuda, France, Germany, Finland, Denmark, Poland, South Africa, Jamaica, Sweden, New Hampshire, New Jersey, New York, Massachusetts, District of Columbia, Illinois, Idaho, Montana, Oregon, and Washington. VII. 2. 1967. (D.A.D.) 1132, 1133. (Both at ISU).

Peniophora pubera (Fries) Sacc.

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from Australia, Austria, Canada, Denmark, France, Germany, Great Britain, Italy, Sweden, Alabama, District of Columbia, Florida, Idaho, Illinois, Kentucky, Louisiana, Maryland, Massachusetts, Missouri, Montana, New Hampshire, New Jersey, New York, Oregon, Rhode Island, Virginia, Washington, and Wisconsin. VII. 2. 1967. (D.A.D.) 1078. (At ISU).

Peniophora burtii Romell

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from New York, New Jersey, Alabama, Louisiana, Ohio, Vermont, West Virginia, Virginia, Illinois, Montana, Massachusetts, and Michigan. VII. 2. 1967. (D.A.D.) 1135, 1136. (Both at ISU).

Sistotrema brinkmanni (Bres.) Erikss.

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from Austria, England, British Guiana, Panama, Nova Scotia, Ontario, Massachusetts, Rhode Island, New York, Virginia, North Carolina, Ohio, Tennessee, Louisiana, Iowa, Idaho, Missouri, and Oregon. VII. 2. 1967. (D.A.D.) 1110, 1068. (Both at ISU).

Corticium furfuraceum Bres.

On decayed deciduous wood. Previously unreported for Pennsylvania. Also known from British Columbia, Ontario, Idaho, North Carolina, and Wisconsin. VII. 2. 1967. (D.A.D.) 1064, 1065. (Both at ISU).

Trechispora farinacea (Pers. ex Fr.) Liberta

On decayed deciduous wood. Reported for Wisconsin and Iowa. Common throughout the eastern United States. VII. 2. 1967. (D.A.D.) 1114, 1080. (Both at ISU).

SUMMARY

Eleven new reports of resupinate Basidiomycetes are noted here for the first time. Although these notes represent no major range extensions, they

aid in filling existing distribution gaps in the eastern United States.

ACKNOWLEDGMENTS

The author wishes to express his thanks to Dr. Anthony E. Liberta and Mr. Robert J. Harris for their ready assistance in confirming and discussing some of the identifications. A special note of thanks is extended to Mr. Michael DeFigio for his help in the collection of these specimens.

LITERATURE CITED

1. HARRIS, R. J. 1965. The Genus *Peniophora* in Illinois. Trans. Ill. Acad. Sci. 59(3).
2. LIBERTA, A. E. 1966. On *Trechispora*. Taxon 15(8):317-319.
3. ———— 1965. Notes on Wisconsin Resupinate Basidiomycetes. Mycologia 57(3):459-464.
4. ———— 1966. Notes on Colorado Resupinate Basidiomycetes. Trans. Ill. Acad. Sci. 59(4).
5. MILLER, L. W. & J. S. BOYLE. 1943. The Hydnaceae of Iowa. University of Iowa Studies in Natural History 18(2).
6. ROGERS, D. P. 1944. The Genera *Trechispora* and *Galzinia* (Thelephoraceae). Mycologia 36(1):70-103.
7. SLYSH, A. R. 1960. The Genus *Peniophora* in New York State and adjacent Regions. State University New York Coll. Forestry Tech. Publ. 83, 95 pages, figs. 1-89.
8. WERESUB, LUELLA K. 1953. Studies of Canadian Thelephoraceae. X. Some species of *Peniophora* Section *Tubuliaferae*. Can. Jour. Bot. 31:760-778.

Manuscript received August 9, 1967.