

## SEPARATION AND DETERMINATION OF THE ALKALI METALS, USING PERCHLORIC ACID

## I. THE SOLUBILITY OF THE PERCHLORATES OF THE ALKALI METALS IN MIXED ORGANIC SOLVENTS

G. FREDERICK SMITH, UNIVERSITY OF ILLINOIS

The quantitative separation and determination of the alkali metals is in general dependent upon the use of organic solvents. The most important separation in this group is that of potassium from sodium. The most common method involves the use of chloroplatinic acid and 80% ethyl alcohol but methods based on the use of perchloric acid have proved satisfactory in this as well as other separations of the alkali metals.<sup>1</sup> Improved methods of procedure in this group depend in part upon a thorough study of new organic solvents.<sup>2</sup> The present work is the first of a series of such studies.

The solubility of the perchlorates of the alkali metals in water and numerous organic solvents, including alcohols and ethyl acetate, has been determined.<sup>2</sup> The nature of the solubility effect obtained using mixed solvents constitutes a marked advantage over the use of individual solvents. The data reported in this paper include the solubility of anhydrous sodium, potassium, rubidium, cesium and ammonium perchlorates as well as sodium perchlorate monohydrate and lithium perchlorate trihydrate in mixtures of methyl, ethyl and N butyl alcohols in various proportions with ethyl acetate. The data are discussed in relation to their application to quantitative analysis.

## INTRODUCTORY CONSIDERATIONS

Quantitative methods for separations in the alkali metal group consist of two types, precipitation and extraction processes. The separation of sodium and lithium by the method of Willard and Smith<sup>1a</sup> is of the former

1a. G. Frederick Smith, J. Am. Chem. Soc., *45*, 2072, (1923).

b. Baxter and Ropert, *ibid.*, *42*, 2046, (1920).

c. Gooch and Blake, Am. J. Sci., *44*, 381, (1917).

d. Morris, Analyst, *45*, 349, (1920). Extensive bibliography.

2A list of organic solvents already studied in this connection includes N. butyl alcohol 1a, also Willard and Smith, J. Am. Chem. Soc., *44*, 2816 (1922); isobutyl alcohol, Winkler, Z. anal. Chem., *52*, 628 (1913); ethyl alcohol containing methyl alcohol 1b; ether and alcohol, Palkin, J. Am. Chem. Soc., *38*, 2326, (1916); amyl alcohol, Gooch, Am. Chem. J., *9*, 33 (1887); and pyridine, Kahlenberg and Krauskopf. J. Am. Chem. Soc., *30*, 1104, (1908).

type. The perchloric acid method for the separation of potassium and sodium is an extraction method. Sodium perchlorate would be more effectively separated from potassium perchlorate by extraction, using an organic solvent, if its solubility could be increased without increasing the solubility of the potassium salt. The selection of a mixture of solvents best suited to this end could be made only after an extended investigation, the results of which have been included in part in this paper.

#### PREPARATION OF MATERIALS

In a former paper<sup>3</sup> the opinion was expressed "that slight amounts of organic impurities, especially those similar in nature to the solvent used, would have no appreciable effect on the solubility results, but even traces of water have a large influence". An alcohol containing a small per cent of an isomeric alcohol or other member of its homologous series would fall under this classification of organic impurities. Such conditions would not hold for some mixtures of organic solvents such as esters contaminated with alcohols or alcohols with aldehydes.

The magnitude of the error resulting from the presence of a very small per cent of dissimilar organic impurity in a particular solvent is shown by the following example:

The solubility of anhydrous sodium perchlorate in a sample of ethyl acetate, physical constants of which duplicated the best values to be found in the literature,<sup>5</sup> was shown to be 8.80% at 25°. The same ethyl acetate containing 0.13% of its volume of anhydrous ethyl alcohol, dissolved 10.74%, an increase of over 22%. From this data it would seem probable that ethyl acetate containing as little as one part of ethyl alcohol in 20000 parts of ethyl acetate would dissolve anhydrous sodium perchlorate to the extent of one per cent in excess of the true value.

As a result of such considerations the solvents used for this work were those of a previous stock which had been carefully purified<sup>3</sup> or carefully selected and compared with the former. Solubility determinations were re-

<sup>3</sup>Willard and Smith, *J. Am. Chem. Soc.*, 45, 286, (1923).

<sup>5</sup>Solvent 13 of reference 3.

peatedly made in establishing the purity of any given solvent. The identity of the samples of the same solvent was established if comparable solubility results were obtained. Solubility data for the pure solvents used are given in reference 3.

*Ethyl Acetate.* Solvent 13, Table 1, reference 3 was used in the case of all solubility determinations except those of potassium, rubidium and cesium perchlorates. These were determined, using a very pure anhydrous product obtainable in quantity from the usual sources. This material without further treatment was found to dissolve 9.11% of anhydrous sodium perchlorate at 25°. This corresponds to a purity of 99.98% assuming the only impurity to be ethyl alcohol.

*Methyl Alcohol.* An anhydrous material purchased in quantity from the usual sources was found to correspond with solvent number 2 of Table 1, reference 3. It was found to dissolve 6.42% of ammonium perchlorate and 34.33% of anhydrous sodium perchlorate as compared with 6.41% and 33.93% for the pure solvent. Technical methyl alcohol, dried over stick potassium hydroxide and distilled, dissolved 42.2% of anhydrous sodium perchlorate.

*Ethyl Alcohol.* 93% alcohol dehydrated with lime to 99.6% was made anhydrous by reaction with a slight excess of metallic calcium. Its boiling range was 78.29°-78.31° corrected, and its density was 0.78515 at 25°/4° corrected to vacuum. The undried alcohol used was found by density determination to be 93.1% in strength.

*N Butyl Alcohol.* The same as solvent number 10, Table 1 of Reference 3.

*Perchloric Acid.* Prepared as given in reference 1a.

*Lithium, Sodium, Ammonium, Potassium, Rubidium and Cesium Perchlorates.* The same materials described in reference 3. The latter three slightly soluble perchlorates were recrystallized from their solution in perchloric acid at the boiling point by cooling to 0° with the object of obtaining fine crystals which required no grinding before testing their solubility. The crystals thus obtained were known to have occluded a small amount of perchloric acid. This fact did not alter solu-



bility data in the organic solvents used, as in no case were the resulting solutions acid. This phenomenon of acid occlusion as exhibited by potassium, rubidium and cesium perchlorates is the subject of the second paper of this series.

*Lithium Perchlorate Trihydrate.* The same material used in reference 3.

*Sodium Perchlorate Monohydrate.* Prepared by crystallization of sodium perchlorate from water below 50°. The material thus obtained was drained centrifugally and dried at room temperature in contact with excess of anhydrous sodium perchlorate. Analysis of the product gave 12.87% water of crystallization as compared to a theoretical value of 12.82%.

#### APPARATUS AND METHODS

The experimental manipulations were carried out essentially as reported in reference 3, pages 290-293 inclusive. All determinations involving potassium, rubidium and cesium perchlorate were carried out, using from 80-100 cc. of the saturated solution in question with conversion of the dissolved perchlorates to sulfates as a means of determination. All determinations were made in duplicate and the strict concordance of the values obtained with those of previously reported determinations<sup>3</sup> is significant.

#### SOLUBILITY DATA

The solubilities in per cent by weight of anhydrous sodium perchlorate, hydrated sodium and lithium perchlorates and ammonium perchlorate in the various mixtures of solvents are listed in Table I.

TABLE I.

SOLUBILITIES OF SODIUM, LITHIUM AND AMMONIUM PERCHLORATES IN ETHYL ACETATE CONTAINING VARIOUS PROPORTIONS OF METHYL AND N-BUTYL ALCOHOLS AT  $25^{\circ} \pm 0.1^{\circ}$

Vol.-% of alcohol	0	5	10	20	30	40	
Vol.-% of ethyl acetate	100	95	90	80	70	60	
Solute used							
NaClO <sub>4</sub> Methyl	8.80	.....	19.39	23.37	25.85	27.81	
NaClO <sub>4</sub> Abs. Ethyl	8.80	13.83	16.05	18.55	19.66	20.08	
NaClO <sub>4</sub> 93% Ethyl	8.80	14.12	16.92	20.00	21.75	22.79	
NaClO <sub>4</sub> N-Butyl	8.80	.....	12.34	13.16	.....	12.82	
NaClO <sub>4</sub> N-Butyl*	8.80	11.97	13.00	14.32	14.75	14.71	
NaClO <sub>4</sub> ·H <sub>2</sub> O Abs. Ethyl	26.32	.....	29.32	31.16	32.44	33.26	
LiClO <sub>4</sub> ·3H <sub>2</sub> O Abs. Ethyl	26.35	.....	31.05	33.59	35.10	36.51	
NH <sub>4</sub> ClO <sub>4</sub> Methyl	0.029	.....	0.135	0.550	1.27	2.31	
Vol. % of alcohol	50	60	70	80	90	95	100
Vol. % of ethyl acetate	50	40	30	20	10	5	0
Solute used							
NaClO <sub>4</sub> Methyl	29.38	30.50	31.61	32.56	33.44	.....	34.33
NaClO <sub>4</sub> Abs. Ethyl	20.07	19.67	18.78	17.53	15.82	14.80	12.83
NaClO <sub>4</sub> 93% Ethyl	23.30	23.39	22.73	21.92	20.96	21.31	20.60
NaClO <sub>4</sub> N-Butyl	11.99	10.54	.....	6.70	4.35	.....	2.19
NaClO <sub>4</sub> N-Butyl*	13.98	13.09	11.66	9.85	7.87	6.97	4.27
NaClO <sub>4</sub> ·H <sub>2</sub> O Abs. Ethyl	33.63	33.61	33.20	32.56	31.56	31.10	29.80
LiClO <sub>4</sub> ·3H <sub>2</sub> O Abs. Ethyl	37.96	38.58	39.79	40.84	41.77	.....	42.16
NH <sub>4</sub> ClO <sub>4</sub> Methyl	3.47	4.35	5.08	5.67	6.12	.....	6.42

\*The same alcohol reported in Ref. 1a; b. p., 112-118°; d<sub>4</sub><sup>25</sup>, 0.8065.

The per cents by volume of the mixed solvents used for the solubilities of each vertical column are given in the first two horizontal rows. The solute used giving the data of each remaining horizontal row is given in the first vertical column. Likewise, the alcohol mixed with the ethyl acetate giving the solubilities contained in each horizontal row is listed in the second vertical column.

The data of Table 1 are represented graphically in Fig. 1, for the purpose of ready comparison.




FIG. I.

The solubility in weight per cent of a given solute is plotted as ordinate. The mixed solvent proportions are plotted as abscissa. The volume per cents of alcohol from 0 to 100 are plotted reading from left to right. Those of ethyl acetate are plotted on the same axis from 0 to 100 reading from right to left.

## DISCUSSION.

From Table 1 and Fig. 1 it is obvious that the solubility of the anhydrous solutes determined are much higher in the mixed solvent than would be predicted from a knowledge of the solubility of the same solute in each individual solvent. Such a predicted series of solubilities would be



represented as a straight line function connecting the first and last point of each curve given. The abnormalities found based on such a consideration range from 100% or even more down.

To select a specific example, consider the solubility of anhydrous sodium perchlorate in a mixture of 20 parts anhydrous ethyl alcohol and 80 parts ethyl acetate. 18.55% by weight of the solution saturated at 25° is found to be sodium perchlorate while 9.6% would be predicted. In this case and that of mixtures of N butyl alcohol with ethyl acetate the individual solvents could be used as the basis of an extraction process for the separation of potassium and sodium, but the advantage of the increased solubility of sodium perchlorate in the mixed solvents is at once apparent. The use of mixed solvents offers no advantage over individual solvents in the case of the extraction of lithium perchlorate.<sup>3</sup> From the data already given, together with the solubility determinations following, a choice of a mixture of solvents to be used for any given separation can be made.

Mixed alcohols give rise to normal solubility effects. This was determined using absolute ethyl alcohol and N butyl alcohol and a straight line solubility curve found. Mixtures of ethyl acetate and acetone give only very slightly abnormal values. The substitution of acetone for ethyl acetate gives only very slightly abnormal solubilities when mixed with alcohols. Mixtures of ethyl acetate with alcohols are thus seen to be the most practicable to employ.

The abnormal solubility results just described could be obtained as a result of two plausible conditions, namely, interaction between the solvents of the mixture, or interaction between the anhydrous solute and one or both solvents. The former condition is indicated through the elimination of the latter as the result of experiment.

Lithium and sodium perchlorates form hydrates. If these solutes in their anhydrous form were dissolved in alcohols, it is possible "alcoholates" instead of hydrates might be formed. In such case the solubility of these modified solutes in alcohols and ethyl acetate might be greater than the anhydrous salts. The abnormal results obtained would thus be explained.

Accordingly the solubility of lithium perchlorate trihydrate in mixtures of ethyl alcohol and ethyl acetates was determined (Curve VII, Figure 1). Its solubility in both solvents is less than that of the anhydrous solute. Only slightly abnormal solubilities were obtained in the mixed solvents. Both circumstances fit the assumptions. With sodium perchlorate monohydrate, however, the solubility in the individual solvents is greater than that of the anhydrous salt and the abnormal solubilities less pronounced (Curve VI, Figure 1).

If the tendency to form "alcoholates" is associated with the formation of hydrates for a given solute, and the former condition gives rise to the abnormal solubility effects obtained in mixed solvents, normal results should be obtained using solutes for which hydrates are not known. The solubility of ammonium perchlorate, representing such an example, in mixtures of methyl alcohol and ethyl acetate was determined (Curve VIII, Figure 1). Here added abnormality was found. For small per cents of alcohol a marked depression of its solubility over the predicted value was obtained. Interaction of solute and solvent is probably therefore not an explanation for the results obtained.

#### THE SOLUBILITY OF POTASSIUM RUBIDIUM AND CESIUM PERCHLORATES IN MIXED ALCOHOLS AND ETHYL ACETATE.

As will be demonstrated in the subsequent papers of this series, the presence of free perchloric acid during the extraction of the soluble from the insoluble alkali perchlorates, using mixed solvents, is a pronounced disadvantage. The addition of perchloric acid for its common ion effect to mixed solvents containing ethyl acetate as the major constituent fails of its purpose. The solubilities considered in this section therefore omit the effect of small amounts of perchloric acid and are reported in Table II.



TABLE II.

The Solubility of Potassium Rubidium and Cesium Perchlorates in Mixed Alcohols and Ethyl Acetate at 25°

Vol. o/o Alcohol		10	15	20
Vol. o/o Ethyl Acetate		90	85	80
Solute	Alcohol used	g./100cc	g./100cc	g./100cc
KClO <sub>4</sub>	Methyl .....	0.0035	0.0052	0.0055
	Ab Ethyl .....	0.0016	0.0026	0.0033
	93 o/o Ethyl.....	0.0028	0.0044	0.0050
	N Butyl .....	0.0024	0.0020	0.0025
RbClO <sub>4</sub>	Methyl .....	0.0024		0.0043
	Ab Ethyl .....	0.0016		0.0010
	93 o/o Ethyl.....			0.0027
	N Butyl .....	0.0011		0.0013
CsClO <sub>4</sub>	Methyl .....	0.0050		0.0056
	Ab Ethyl .....	0.0013		0.0023
	93 o/o Ethyl.....			0.0040
	N Butyl .....	0.0009		0.0012

The description of Table I already given serves likewise for Table II. The values reported in Table II cannot be considered of greater accuracy than approximately 0.5 milligram. All errors accumulating throughout the process of their determination tend to give high results. The data represent saturated solutions, a condition not obtained in practical applications. Predictions based upon the data given in Table II may be somewhat liberally interpreted.

The solubility of potassium perchlorate in pure ethyl acetate is 1.3 milligrams in 100 cc. at 25 (reference 3). Rubidium and cesium perchlorates are insoluble in ethyl acetate. Ethyl acetate should therefore be the wash solution in all cases. If one extraction was found sufficient, 20 cc. of extractant, representing one-fifth the solubilities given in Table II, would give the maximum error through solubility. Working at less than 25°, which is ordinary practice, would decrease this error. Any of the combinations of Table II could then be employed. If two extractions were found necessary, approximately 30 cc. of extractant would be required and the higher values of Table II would represent too great errors.

Absolute ethyl alcohol and ethyl acetate give the best values if the figures of Table I are also considered. 93%

ethyl alcohol can be substituted for absolute alcohol with but small decrease in accuracy. N butyl alcohol and ethyl acetate give the lowest solubilities and could probably be used in all proportions.

The data of the present paper are thus seen to be a practical guide to the selection of most probably favorable conditions for any of the separations involved. Exact choices depend upon a more thorough study of each case.

#### SUMMARY

1. The solubility of the alkali perchlorates in various mixtures of methyl, ethyl and N butyl alcohol with ethyl acetate have been determined.
2. The solubility data thus obtained were shown to represent conditions advantageous in the separation and determination of the alkali metals by the extraction process.
3. An attempt was made to interpret the abnormal solubility results and to point out their analytical significance.
4. Solubility determinations were shown to be of value in establishing the purity of the solvents employed.

Contribution from the  
Chemical Laboratory  
of  
The University of Illinois